


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)


[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)

**Scholar** All articles - **Recent articles** Results **1 - 10** of about **227** for **man glnac antibody thiol OR Cysteine iodoacetamide** with **Safesearch on**. (0.29 seconds)

**Mass Spectrometry of the Humanized Monoclonal Antibody Campath 1H** - [all 3 versions »](#)

DS Ashton, OR Beddell, DJ Cooper, SJ Craig, AC ... - Analytical Chemistry, 1995 - pubs.acs.org

... **DL-Cysteine**, dithiothreitol ... variant of bacterial subtilisin and a bacterial **thiol** oxidoreductase were ... Chester, UK A solution of CAMPATH 1H **antibody** in phosphate ...

[Cited by 18](#) - [Related Articles](#) - [Web Search](#)

**[CITATION] Plant-derived mouse IgG monoclonal antibody fused to KDEL endoplasmic reticulum-retention signal is ...** - [all 2 versions »](#)

A Triguero, G Cabrera, JA Cremata, CT Yuen, J ... - Plant Biotechnology Journal, 2005 - Blackwell Synergy

... C with anti- $\beta$ 1,2-Xyl and anti- $\alpha$ 1,3-Fuc **antibodies** (diluted 1 ... Relative retention times were calculated, referred to that of the standard **Man 3 GlcNAc 2-2AB** ...

[Cited by 18](#) - [Related Articles](#) - [Web Search](#)

**Impact of variable domain glycosylation on antibody clearance: An LC/MS characterization** - [all 3 versions »](#)

L Huang, S Biotsi, KR Bales, U Kuchibhotla - Analytical Biochemistry, 2006 - Elsevier

... Another **GlcNAc** must be linked to only available HexNAc ... its linkage to GlcNAc $\beta$  (1  $\rightarrow$  2)**Man** could not ... captured using immobilized goat anti-human Fc $\gamma$ **antibody** ...

[Cited by 11](#) - [Related Articles](#) - [Web Search](#)

**[CITATION] Structural Characterization of a Recombinant Antibody Produced in Yeast**

PC Gough, IA Kadura, GM Kelly, A Kreuzman, J Larson ...

[Related Articles](#) - [Web Search](#)

**Biochemical and functional characterization of the Tn-specific lectin from Salvia solarea seeds** - [all 4 versions »](#)

A Medeiros, S Bianchi, JJ Calvete, H Baker, S Bay ... - FEBS Journal, 2000 - FEBS

... Anti-Tn monoclonal **antibody** 83D4 (IgM, ) [22] was ... molar excess of **iodoacetamide** over **cysteine** residues; and ... chains of SSL are composed of **GlcNAc**, **Man**, **Xyl**, and ...

[Cited by 17](#) - [Related Articles](#) - [Web Search](#) - [BI Direct](#)

**N-Glycosylation of a mouse IgG expressed in transgenic tobacco plants** - [all 3 versions »](#)

M Cabanes-Macheteau - Glycobiology, 1999 - Soc Glycobiology

... **GlcNAc 2** core substituted by terminal **GlcNAc** residues ... **Man-8**), and soybean agglutinin (**Man-9**) as ... to pulsed amperometric detection; MA, monoclonal **antibody**; 1 H ...

[Cited by 119](#) - [Related Articles](#) - [Web Search](#) - [BI Direct](#)

**... Specific Glycosylation of an Aglycosylated Human IgG1-Fc Antibody Protein Generates Neoglycoproteins ...** - [all 11 versions »](#)

GM Watt, J Lund, M Levens, VSK Kottli, R Jefferis, ... - Chemistry & Biology, 2003 - Elsevier

... 1). Epitope mapping using an exten- sive panel of monoclonal **antibodies** directed

against the human IgG1 **antibody** revealed that ... [-S(**GlcNAc**) 2 **Man**] 15 25,030 ...

[Cited by 11](#) - [Related Articles](#) - [Web Search](#)

Glycosylation of the variable region of immunoglobulin G—site specific maturation of the sugar ... - all 4 versions »

T Erdo, A Wright, SL Morrison, A Kobata - Molecular immunology, 1995 - Elsevier

... Purified secreted **antibody** from each transfectoma ... M, EDTA to 2 mM and **cysteine** to

10 ... **Man**?1 + 4**GlcNAc**Bl+ 4**GlcNAc**OT (**Man** - **GlcNAc** - **GlcNAc**,) was prepared from ...

[Cited by 15](#) - [Related Articles](#) - [Web Search](#) - [BI Direct](#)

The cattle tick antigen, Bm95, expressed in Pichia pastoris contains short chains of N- and O-glycans - all 4 versions »

LJ González, JA Cremata, Y Guanche, Y Ramos, A ... - Archives of Biochemistry and Biophysics, 2004 - Elsevier

... 7]. However, B. microplus isolations show low susceptibility to **antibodies** developed

in ... a database of the retention times relative to **Man** 7 **GlcNAc** 2 -4ABA ...

[Cited by 3](#) - [Related Articles](#) - [Web Search](#)

Mass spectrometric analysis reveals a **cysteine** bridge between residues 2 and 61 of the auxin-binding ... - all 3 versions »

C Feckler, G Muster, W Feser, A Römer, K Palme - FEBS Letters, 2001 - Elsevier

... were based on epitope mapping using monoclonal **antibodies** against peptides ... type sugar

predicted from earlier results was considered (**GlcNAc** 2 **Man** 9 , 1865.6 ...

[Cited by 4](#) - [Related Articles](#) - [Web Search](#)

Key authors: [D Williams](#) - [A Helenius](#) - [A Parodi](#) - [E Trombetta](#) - [B Davis](#)

Google

Result Page: 1 2 3 4 5 6 7 8 9 10 [Next](#)

man glnac antibody thiol OR Cyste

Search

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google